


Idaho: Most Common Causes of Death

- Total cardiovascular diseases (principally ischemic heart disease and stroke) are the most common cause of death in Idaho, all cancers are the second most common cause, chronic obstructive pulmonary disease is fourth, and diabetes is seventh.
- In 1995, 70% of all deaths in Idaho were due to the four most common chronic disease causes of death.
- The death rate for chronic obstructive pulmonary disease was higher in Idaho than in the United States; the death rates for total cardiovascular diseases, all cancers, and diabetes were lower.

Causes of Death, Idaho Compared With United States, 1995*




*All data are age adjusted, 1970 total U.S. population.

**Total cardiovascular disease death rates include rates of death due to ischemic heart disease (109.8 per 100,000 in Idaho and 135.2 per 100,000 in the United States) and rates of death due to stroke (38.9 per 100,000 in Idaho and 42.5 per 100,000 in the United States).

Idaho: Cardiovascular Diseases


- Total cardiovascular diseases are the most common cause of death in Idaho, accounting for 36% of all deaths.
- Ischemic heart disease accounted for 51% of all cardiovascular disease deaths in Idaho in 1995; 1,625 people in Idaho died of ischemic heart disease.
- In 1995, 637 people in Idaho died of stroke.

Idaho: Total Cardiovascular Disease Death Rates
By Sex and Race, 1995




**Too few numbers to analyze.

Idaho: Ischemic Heart Disease Death Rates
By Sex and Race, 1995



**Too few numbers to analyze.

Idaho: Stroke Death Rates
By Sex and Race, 1995




**Too few numbers to analyze.

Note: All data are age adjusted, 1970 total U.S. population.

Idaho: Cancer


- Cancer accounted for 25% of all deaths in Idaho in 1995; 2,006 people in Idaho died of cancer.
- In Idaho in 1995, 482 people died of lung cancer, 180 people died of colorectal cancer, and 157 women died of breast cancer.
- The American Cancer Society estimates that 570 new cases of lung cancer, 450 new cases of colorectal cancer, and 720 new cases of breast cancer will be diagnosed in Idaho in 1997.

Idaho: All Cancer Death Rates
By Sex and Race, 1995




**Too few numbers to analyze.

Idaho: Lung Cancer Death Rates
By Sex and Race, 1995




**Too few numbers to analyze.

Idaho: Colorectal Cancer Death Rates
By Sex and Race, 1995



**Too few numbers to analyze.

Idaho: Breast Cancer Death Rates Among Women
By Race, 1995




**Too few numbers to analyze.

Note: All data are age adjusted, 1970 total U.S. population.

Idaho: Diabetes

- In 1994, 29,894 adults in Idaho had diagnosed diabetes.
- Diabetes was the underlying cause of 206 deaths in Idaho in 1995.
- In 1993, diabetes was the most common contributing cause of 48 new cases of end-stage kidney disease in Idaho.
- Diabetes is believed to be underreported on death certificates, both as a condition and as a cause of death.




Note: All data are age adjusted, 1970 total U.S. population.

Idaho: Risk Factors

Idaho: Percentage of Adults Who Reported Cigarette Smoking*

By Sex, Race, and Hispanic Origin, 1996 BRFSS




*Ever smoked at least 100 cigarettes and now smoke every day or some days.

**Too few numbers to analyze.

Idaho: Percentage of High School Students Who Reported Cigarette Smoking*


By Sex, 1995 YRBSS



*Smoked cigarettes on 1 or more of the 30 days preceding the survey.

Idaho: Percentage of Adults Who Reported No Leisure-Time Physical Activity*

By Sex, Race, and Hispanic Origin, 1996 BRFSS




*No exercise, recreation, or physical activity (other than regular job duties) during the previous month.

**Too few numbers to analyze.


Idaho: Percentage of High School Students Not Enrolled in Physical Education Classes

By Sex, 1995 YRBSS



Idaho: Percentage of Adults Who Reported Being Overweight*

By Sex, Race, and Hispanic Origin, 1996 BRFSS




*Body mass index ≥ 27.8 kg/m² for men and ≥ 27.3 kg/m² for women.


**Too few numbers to analyze.

Note: All BRFSS data are provisional and age adjusted, 1970 total U.S. population. YRBSS data are unweighted and thus apply only to the students participating in the survey.


Idaho: Risk Factors




**Too few numbers to analyze.



Idaho: Preventive Services



**Too few numbers to analyze.



**Too few numbers to analyze.

Note: All BRFSS data are provisional and age adjusted, 1970 total U.S. population. YRBSS data are unweighted and thus apply only to students participating in the survey.